# Frontline First Aid **Lesson Plan Example – Scenario Presentation**



Call	Detai	ls:

**Unresponsive Patient** 

Instructions to role playe
----------------------------

Patient is unresponsive for the entire call. Bystander can answer all pertinent questions.

Participant's name: _	Time Started:	

	Critical Findings	Critical Actions or Interventions	Yes	No	Notes and Comments			
Scene Survey								
Hazards	None	Gloves, Goggles						
MOI	Gently laid down							
# of Patients	1							
Additional Resources Needed								
		Approach	•					
Identify self		Introduce yourself						
What Happened?	Laid down and passed out	Talk to the bystander						
Please do not move (if applicable)	N/App							
		Primary Survey						
C-Spine (if necessary)	Not needed							
LOC (AVPU)	Unresponsive							
Airway	Clear	OPA						
Breathing	12; effective							



	Critical Findings	Critical Actions or Interventions	Yes	No	Notes and Comments
Circulation		interventions			
Skin	Pale, Cool, Clammy				
Pulse Check	Present				
RBS	No blood				
Critical Interventions					
Oxygen		10 lpm; Simple Mask			
Blanket		Cover with blanket			
Bleeding					
Position		Keep comfortable			
Pulse Oximeter	97	Apply pulse oximeter			Recorded with Vitals
Transport Decision	☐ Rapid Transport (Unresponsive)	Partner prepares equipment			Continue protocols until transport
		Secondary Survey			
History					
<b>S</b> igns and Symptoms	Dizziness				
Allergies	Penicillin				
<b>M</b> edications	none				
Past medical Hx	none				
Last Meal	½ hour ago				
Events leading up	Sitting at home				
<b>O</b> nset	Sudden				
Provoke	Standing				
Quality	Dizziness				
<b>R</b> egion/Radiate	Head				
<b>S</b> everity	6/10				
Time	8 minutes ago				



	Critical Findings	Critical Actions or Interventions	Yes	No	Notes and Comments
Vital Signs					Second set of Vitals
Α					
V	Unresponsive				Unresponsive
Р	'				'
U					
Respirations					
Rate	12				12
Rhythm	Regular				Regular
Quality	Effective				Effective
Pulse					
Rate	70				80
Rhythm	Regular				Regular
Quality	Strong				Strong
Skin Condition	Pale; Cool, Sweaty				Pale; Cool, Sweaty
Blood Pressure	110/60				115/70
Pupils	3 mm; Equal; Reactive				3 mm; Equal; Reactive
Oxygen Saturation	Palpable pulse is within 10 bpm of Pulse Oximeter	Simple Mask; 10 lpm  ☐ Compare Oximeter pulse rate to palpated rate ☐ Difference < 10 bpm considered accurate			98%
If Inaccurate	□ Difference greater than 10 bpm	<ul> <li>□ Remove from finger</li> <li>□ Use warmer finger</li> <li>□ Remove nail polish</li> <li>□ Use Toe or earlobe if appropriate</li> <li>□ Re-apply oximeter and compare</li> </ul>			



	Critical Findings	Critical Actions or Interventions	Yes	No	Notes and Comments
Head to Toe					
Head					
Look	Unremarkable				
Feel	Unremarkable				
Neck					
Look	Unremarkable				
Feel	Unremarkable				
Chest					
Look	Unremarkable				
Listen/Auscultate	Equal-Bilateral				
Feel	Unremarkable				
Abdomen					
Look	Unremarkable				
Feel	Unremarkable				
Pelvis					
Look	Unremarkable				
Feel	Unremarkable				
Lower Extremities					
Look	Unremarkable				
Feel	Unremarkable				
Pulse	Present				
Motor	Normal				
Sensory	Normal				
Upper Extremities					
Look	Unremarkable				
Feel	Unremarkable				
Pulse	Present				
Motor	Normal				
Sensory	Normal				
Back					
Look	Unremarkable				
Feel	Unremarkable				
	Critical Findings	Critical Actions or	Yes	No	Notes and Comments
		Interventions			



	Critical Findings		Critical Actions or Interventions		Yes	No	Notes and Comments	
Monitor								
Oxygen Saturation	98%							Treat the patientNOT the oximeter
Appropriate Treatment	More advanced care needed		☐ Check Vitals every 5 minutes					
Time Ended		Succes	sful	Yes	No			
Comments:								